

The Formation of Disks in Elliptical Galaxies

Thorsten Naab & Andreas Burkert

Max-Planck-Institut für Astronomie, Königstuhl 17,

D-69117 Heidelberg,

Germany

Received _____; accepted _____

ABSTRACT

We investigate detailed kinematical properties of simulated collisionless merger remnants of disk galaxies with mass ratios of 1:1 and 3:1. The simulations were performed by direct summation using the new special hardware device GRAPE-5. In agreement with observations, the shape of the line-of-sight velocity distribution (LOSVD) is Gaussian with small deviations. For most cases we find that the retrograde wings of the LOSVD are steeper than the prograde ones. This is in contradiction with observations which show broad retrograde and steep prograde wings. This serious problem in the collisionless formation scenario of massive elliptical galaxies can be solved if all rotating ellipticals, even boxy ones, contain an additional stellar disk component with $\propto 15\%$ of the total stellar mass and a scale length of order the effective radius of the spheroid. We propose that the progenitor galaxies of massive ellipticals must have contained a significant amount of gas that did not condense into stars during the merger process itself but formed an extended gaseous disk before the star formation epoch. The heating source that prevented the gas from forming stars early and the origin of the large specific angular momentum required for the gas component to form an extended disk are still unsolved problems.

Subject headings: galaxies: elliptical – galaxies: interaction – galaxies: structure – galaxies: evolution – methods: numerical

1. Introduction

The merger hypothesis (Toomre & Toomre 1972) assumes that elliptical galaxies form by major mergers of spiral galaxies. This is supported by observations of nearby merging

systems that seem to evolve into systems with properties comparable to elliptical galaxies (Hibbard & Yun 1999). The gravitational interaction during the merger process also triggers massive star formation and, probably, the formation of massive central black holes and AGNs (Sanders & Mirabel 1996; Rigopoulou et al. 1999). On the theoretical side, numerous numerical simulations have shown that mergers lead to spheroidal systems with global properties comparable to elliptical galaxies (Barnes & Hernquist 1992; Barnes 1998; Bekki & Shioya 1997; Naab, Burkert & Hernquist 1999). During the merging epoch the galaxies are far from dynamical equilibrium. Gas, if added to the simulations, condenses into stars by massive bursts of star formation during the merging process or effectively loses angular momentum and falls into the central parts of the merger remnant, probably feeding a central black hole (Mihos & Hernquist 1996; Springel 2000; Bekki 2000). When the merger remnants have settled into dynamical equilibrium, phase mixing and violent relaxation have erased most information about the initial conditions. However, as violent relaxation is incomplete, the stellar phase space distribution of elliptical galaxies should still show fine structure which affects the isophotal shapes and velocity distributions. This fine structure provides fundamental insight into the formation epoch of elliptical galaxies.

Numerical simulations of collisionless mergers of disk galaxies have confirmed this conclusion. The LOSVDs of merger remnants and the orbit structure of their stars contain information about the initial disk orientations (Heyl, Hernquist & Spergel 1996). The isophotal shapes and the dynamics of the merger remnants are primarily determined by their initial mass ratios (Naab, Burkert & Hernquist 1999; Bendo & Barnes 2000).

In this letter we show that the LOSVDs of most elliptical galaxies contain evidence for a second disk component with roughly 10% to 20% the luminosity of the spheroid and a large scale radius which is of order the effective radius of the spheroid. This disk might have formed by late gas infall from gas-rich progenitors, followed by star formation.

Observationally, photometric and kinematic investigations of elliptical galaxies have already indicated the existence of large stellar disks of the same order of magnitude as predicted here, especially in disk, rotating ellipticals (Rix & White 1990; Scorza et al. 1998).

We demonstrate that this component is required in almost all ellipticals, independent of whether they are disk or boxy in order to bring the observations in agreement with the merger scenario of the formation of early-type galaxies.

2. The merger models

We use the method described by Hernquist (1993) to construct disk galaxies in dynamical equilibrium. The system of units is: gravitational constant $G=1$, exponential scale length of the larger disk in the merger $h=1$ and mass of the larger disk $M_d = 1$. The disks are exponential with an additional spherical, non-rotating bulge with mass $M_b = 1/3$, a Hernquist density profile (Hernquist 1990) and a scale length $r_b = 0.2h$. The system lives in a spherical pseudo-isothermal halo with a mass $M_d = 5.8$, cut-off radius $r_c = 10h$ and core radius $\gamma = 1h$.

The N-body simulations were performed using a direct summation code with the new special purpose hardware GRAPE-5 (Kawai et al. 2000). The 1:1 merger was calculated adopting in total 400000 particles with each galaxy consisting of 60000 disk particles, 20000 disk particles and 120000 halo particles. For the 3:1 merger the parameters of the more massive galaxy were as described above. The low-mass galaxy contained 1/3 the mass and number of particles of the larger galaxy, with a disk scale length of $h = \sqrt{1/3}$, as expected from the Tully-Fisher relation.

For both mass ratios, the galaxies approached each other on nearly parabolic orbits with an initial separation of 30 length units and a pericenter distance of 2 length units. In

this letter we focus on one corotating and one counterrotating geometry. We have found that these two impact parameters are representative for a much larger set of simulations that have been performed and that will be discussed in a subsequent paper. The inclinations of the corotating disk models (called model A for the equal mass merger and model C for the 3:1 merger) relative to the orbital plane are $t_1 = 30^\circ$ and $t_2 = -30^\circ$ with arguments of pericenter of $\omega_1 = 30^\circ$ and $\omega_2 = -30^\circ$, respectively. The counterrotating disks (called model B for 1:1 and model D for 3:1) have $t_1 = 30^\circ$ and $t_2 = 150^\circ$, $\omega_1 = 30^\circ$ and $\omega_2 = -30^\circ$, respectively. In all simulations the merger remnants were allowed to settle into dynamical equilibrium for approximately 10 dynamical timescales after the merger was complete. Then their equilibrium state was analysed.

3. LOSVD analysis of the merger remnants

To measure the line-of-sight velocity distribution of a merger remnant we shifted the densest region of every two-dimensional projection to the origin. Then we placed a slit with a width of 0.2 unit lengths along the apparent long axis of each projected remnant. Thereafter we binned all particles falling within each grid cell in velocity along the line-of-sight. The grid spacing was chosen to be 0.15 times the projected half-mass radius. The width of the velocity bins was set to a value of 0.2 for line-of-sight velocities v_{los} in the range $-4 \geq v_{\text{los}} \geq 4$. This results in 80 velocity bins over the whole velocity interval. Using the binned velocity data we constructed line-of-sight velocity profiles (LOSVD) for each bin along the grid. Subsequently we parametrized deviations from the Gaussian shape of the velocity profile using Gauss-Hermite basis functions (van der Marel & Franx 1993; Gerhard 1993; Bendo & Barnes 2000). The kinematic parameters of each profile (σ_{fit} , v_{fit} , H_3 , H_4) were then determined by least squares fitting. The large number of simulated stellar particles (> 100000) guaranteed that at least 2000 particles fall within each slit

inside one effective radius.

4. Comparison with observations

The straight lines in Figure 1 indicate the observed local correlation between H_3 and v/σ for a sample of elliptical galaxies in low density environments, published by Bender, Saglia & Gerhard (1994). In all cases H_3 and v/σ have opposite signs. The data indicate that all elliptical galaxies have LOSVDs with steep prograde wings and broad retrograde wings. Mehlert et al. (2000) have shown that this result also holds for cluster ellipticals in Coma.

The local correlation between H_3 and v/σ for the simulated corotating (A,C) and counterrotating (B,D) 1:1 and 3:1 mergers, respectively, is shown by the dots in Figure 1. Every remnant is analysed as seen from 50 random viewing angles. For the corotating mergers the correlation between H_3 and v/σ is almost opposite to the observed one. This is also reflected in a positive effective value for H_3 (left plot in Figure 2) for almost all projection angles, where $\langle H_3 \rangle$ is defined as the mean value between 0.25 and 0.75 effective radii (Bender, Saglia & Gerhard 1994). Here, a positive $\langle H_3 \rangle$ corresponds to an anticorrelation between H_3 and v/σ , such that $H_3 > 0$ for $v/\sigma < 0$ and vice versa. We find that $\langle H_3 \rangle$ versus v/σ_0 does not follow the observed correlation, indicated by the solid and dashed lines. The profiles have broad prograde wings and narrow retrograde wings. The counterrotating 3:1 merger (D) in Fig. 1 shows a large spread around zero but also does not agree with observations. The only exception is the equal mass merger of counter-rotating disks (B) which leads to a very anisotropic elliptical with no signature of rotation.

Note, that the four simulations which we focus on in this paper have been chosen because they are representative for a much larger set of simulations with different orbital

geometries. These four models have been recalculated with high resolution for better statistics.

We conclude that collisionless mergers of disk galaxies in general fail to explain the detailed kinematics of all observed elliptical galaxies (also massive, boxy ones) that exhibit a significant amount of rotation ($v/\sigma \geq 0.2$) inside one r_{eff} . Further evidence for a possible failure of the collisionless merger picture comes from Cretton et al. (2001) who showed that the kinematical properties of very faint and fast rotating, disk elliptical galaxies can not be explained by collisionless mergers of disk galaxies.

5. Theoretical evidence for disks in elliptical galaxies?

One plausible explanation for line-of-sight velocity distributions with negative H_3 is the superposition of a spheroidal body with a disk-like component (Bender, Saglia & Gerhard 1994). With a simple experiment one can test if the wrong correlations found in our remnants result from a lack of such a component. We artificially added a thin (scale height $\approx 0.05r_{\text{eff}}$), cold (velocity dispersion perpendicular to the disk $\sigma_z = 0$), stellar disk with an exponential surface density profile. The disk was placed in the plane defined by the long and intermediate axes of the main stellar body, rotating in the same direction as the main stellar body. The particles of the disk were assumed to move around the center of the galaxy in centrifugal equilibrium with the gravitational potential arising from the total enclosed mass. No additional random motion was added. Under these simple assumptions only two parameters remain free: the total mass of the disk M_d in units of the total luminous mass of the remnant and its scale length r_d in units of the projected half-light radius r_{eff} . The results for the corotating 3:1 remnant (C) are summarized in the Figures 3 and 4. This case is representative for all our unequal mass merger remnants. In general, we find that disks with small masses or radii do not change the LOSVDs of the stellar component. Disks with

masses and radii in the region indicated by the black dots in Figure 4, on the other hand, lead to a significant change in the resulting line profile. The prograde wings become steeper than the retrograde ones in very good agreement with observations. This effect is shown in Figure 3 for a disk with 15% of the spheroid mass and $r_d = 1.25r_{\text{eff}}$. The influence on the $\langle H_3 \rangle$ versus v/σ_0 correlation and its dependence on projection effects is shown on the right hand side of Figure 2. Now the values fall in the observed regime.

If the additional disk becomes too massive, the absolute values for H_3 are larger than observed and the surface brightness profiles change from de Vaucouleurs to exponential profiles which is again not in agreement with observations. We therefore conclude that the existence of an additional stellar disk component with 10% to 20% the luminosity of the spheroid and a scale length of the order of r_{eff} can explain the observed correlation between H_3 and v/σ in elliptical galaxies.

6. Discussion and Conclusions

Our simulations indicate that models of pure collisionless mergers in general fail to reproduce LOSVDs of observed elliptical galaxies. All simulated profiles locally have the wrong sign of H_3 with respect to (v/σ) , compared with the observations. A similar result has been reported by Bendo & Barnes (2000). They, however, did not investigate projection effects. A detailed investigation of local properties of the LOSVDs and projection effects shows that the presence of an additional stellar disk component added to the stellar body after the merger is complete could solve the problem.

One possible origin for such a disk is gas that must have been present during the formation epoch of massive ellipticals and that formed a disk after the major merger event was completed. In order to fit the observed profiles, the amount of gas that settled into

the equatorial plane must have been significant. It is puzzling why the gas did not turn into stars prior to its infall into the equatorial plane and how the gas could keep its angular momentum required for the disk to have a scale length equal to the effective radius of the spheroid.

Barnes & Hernquist (1996) do find gaseous disk-like components in their equal mass merger models. Tidal torques lead however to efficient angular momentum loss in the gaseous component resulting in gas infall to the center. Only 20% of the initial gas mass (which was 10% of the initial disk mass) and by this less than 2% of the stellar mass settled into an extended disk-like component. In this case the disk would not be massive enough to change the LOSVD of the system significantly. Equal mass mergers of very gas rich disks (more than 50% gas) could, however, lead to the formation of more massive disk-like components in the end.

First simulations by Naab, & Burkert (2001) have shown that for unequal mass mergers of gas rich galaxies which lead to disky, fast rotating ellipticals due to the stronger centrifugal support, a large fraction of the gas settles into extended disks after the merger is complete, if star formation is suppressed. This seems to be an attractive scenario to explain the existence of large disks in disky elliptical galaxies. The origin of large stellar disks in massive boxy ellipticals with a small but significant amount of rotation that form presumably from equal-mass mergers is, however, still unclear.

We thank Ralf Bender, Hans-Walter Rix and Nicola Cretton for helpful discussions.

REFERENCES

- Barnes, J. E. & Hernquist, L. 1992, ARA&A, 30, 705
- Barnes, J. E. & Hernquist, L. 1996, ApJ, 471, 115
- Barnes, J., Galaxies: interactions and induced star formation: lecture notes 1996 / Saas Fee
Advanced Course 26, eds. D. Friedli, L. Martinet, and D. Pfenniger, Springer, 1998
- Bendo, G. J. & Barnes, J. E. 2000, MNRAS, 316, 315
- Bender, R., Saglia, R. P., & Gerhard, O. E. 1994, MNRAS, 269, 785
- Bekki, K. & Shioya, Y. 1997, ApJ, 478, L17
- Bekki, K. 2000, ApJ, 545, 753
- Cretton, N., Naab, T., Rix, H.-W., Burkert, A. 2001, accepted by ApJ
- Gerhard, O. E. 1993, MNRAS, 265, 213
- Hernquist, L. 1990, ApJ, 356, 359
- Hernquist, L. 1993, ApJS, 86, 389
- Heyl, J. S., Hernquist, L. & Spergel, D. N. 1996, ApJ, 463, 69
- Hibbard, J. E. & Yun, M. S. 1999, ApJ, 522, L93
- Kawai, A., Fukushige, T., Makino, J. & Taiji, M. 2000, PASJ, 52, 659
- Mehlert, D., Saglia, R. P., Bender, R., & Wegner, G. 2000, A&AS, 141, 449
- Mihos, J. C. & Hernquist, L. 1996, ApJ, 464, 641
- Naab, T., Burkert, A., & Hernquist, L. 1999, ApJ, 523, L133

- Naab, T., & Burkert, A., 2000, to appear in ASP Conf. Ser. "Galaxy Disk and Disk Galaxies", Eds. J.G. Funes S.J. and E.M. Corsini
- van der Marel, R. P. & Franx, M. 1993, ApJ, 407, 525
- Rigopoulou, D., Spoon, H. W. W., Genzel, R., Lutz, D., Moorwood, A. F. M., & Tran, Q. D. 1999, AJ, 118, 2625
- Rix, H. & White, S. D. M. 1990, ApJ, 362, 52
- Scorza, C., Bender, R., Winkelmann, C., Capaccioli, M., & Macchetto, D. F. 1998, A&AS, 131, 265
- Sanders, D. B. & Mirabel, I. F. 1996, ARA&A, 34, 749
- Springel, V. 2000, MNRAS, 312, 859
- Toomre, A. & Toomre, J. 1972, ApJ, 178, 623




Fig. 1.— Local correlation between H_3 and v/σ for the 1:1 and 3:1 merger simulations. The models (A,C) and (B,D) have co -and counterrotating geometries, respectively. The dots represent the data for each model as seen from 50 random viewing angles. The error bar in the upper panel shows the bootstrap error at $0.75 r_{\text{eff}}$. The observed correlation from Bender, Saglia & Gerhard (1994) is tentatively indicated by the straight lines. For the complete data set see their Figure 15.




Fig. 2.— Left: Effective H_3 versus v/σ_0 (rotational velocity at one r_{eff} over central velocity dispersion) for 50 random projection of model (C). Right: Same for model (C) with an additional disk as described in the text. The straight line is the mean correlation observed by Bender, Saglia & Gerhard (1994). The parallel long-dashed lines give an indication of the observed spread. The short dashed line is $\langle H_3 \rangle = 0$.




Fig. 3.— Local correlation between H_3 and v/σ for model (C) with an additional thin stellar exponential disk with a scale length of $1.25r_{\text{eff}}$. A mass of 15% of the spheroid is assumed. The error is derived in the same way as in Figure 1.




Fig. 4.— Mass M_d versus scale radius r_d for a disk added to the merger remnant (C). Combinations that are able to reproduce (big dot), almost reproduce (small dot) or fail to reproduce (minus sign) the observed correlations are shown.